

# GPDs and DVCS with Positrons

Matthias Burkardt

*Department of Physics, New Mexico State University, Las Cruces, NM 88003, and  
Jefferson Lab, 12000 Jefferson Ave., Newport News, VA 23606.*

**Abstract.** The beam charge asymmetry helps to isolate the real part of the deeply virtual Compton scattering (DVCS) amplitude. It is discussed what information can be gained both from the real and imaginary part of the DVCS amplitude.

## INTRODUCTION

Form factors are the coherent amplitude that the nucleon remains intact when one of its quarks absorbs a certain momentum transfer  $\Delta^\mu$ . In this amplitude, the contribution from all quarks carrying all kinds of momenta is added up coherently. Compton scattering provides a more surgical approach (Fig. 1). Due to the presence of the quark propagator between the two photon vertices (note that for virtual photons with large virtuality, the Compton amplitude is dominated by handbag diagrams), the Compton amplitude is sensitive to the momentum fraction  $x$  carried by the active quark.

Suppose one could dissect the Dirac form factor  $F_1^q$  or the Pauli form factor  $F_2^q$  for quarks with flavor  $q$  w.r.t. the (average) momentum fraction  $x = \frac{1}{2}(x_i + x_f)$  carried by the active quark. The result would be the generalized parton distributions  $H^q(x, \xi, t)$  and  $E^q(x, \xi, t)$  respectively

$$F_1^q(t) = \int_{-1}^1 dx H^q(x, \xi, t) \quad F_2^q(t) = \int_{-1}^1 dx E^q(x, \xi, t), \quad (1)$$

where  $t = \Delta^2$ . As the variable  $x$  represents the (average) momentum fraction of the active quark in the ‘infinite momentum’ direction, it makes a difference whether the momentum transfer is parallel or perpendicular to that direction. This ‘angular dependence’ is provided by the dependence of GPDs on the variable  $\xi = \frac{1}{2}(x_f - x_i)$ . Since the information about the infinite momentum direction is ‘lost’ when the quark momentum is not specified, the  $\xi$ -dependence of GPDs must disappear when  $x$  is integrated over, as in (1). Also a consequence of Lorentz invariance, a whole set of ‘polynomiality conditions’ exists for higher moments of GPDs. For example, the  $x^{n-1}$ -moments of GPDs, with  $n$  even, are even polynomials in  $\xi$  with highest power equal to  $n$

$$\int_{-1}^1 dx x^{n-1} H^q(x, \xi, t) = A_{n0}^q(t) + A_{n2}^q(t) \xi^2 + \dots + A_{nn}^q(t) \xi^n, \quad (2)$$

and similar for  $E^q(x, \xi, t)$ . These conditions provide highly nontrivial constraints for the  $x, \xi$ -dependence and may play a crucial role in the GPD extraction from the DVCS amplitude.




**FIGURE 1.** Comparison between Compton scattering (a) and form factors (b). The Compton amplitude (handbag diagrams dominate for large virtualities of the virtual photon), involves a quark propagator between the two photon vertices




**FIGURE 2.** Interference between the DVCS amplitude (a) and the Bethe-Heitler process (b,c).

GPDs enter the DVCS amplitude  $\mathcal{A}_{DVCS}$  through convolution integrals

$$\mathcal{A}_{DVCS} \sim \int_{-1}^1 dx \frac{GPD(x, \xi, t)}{x - \xi + i\epsilon}, \quad (3)$$

i.e. the imaginary part of the DVCS amplitude depends only on GPDs along the ‘diagonal’  $x = \xi$

$$\Im \mathcal{A}_{DVCS}(\xi, t) \sim GPD(\xi, \xi, t), \quad (4)$$

while the real part also probes GPDs for  $x \neq \xi$

$$\Re \mathcal{A}_{DVCS}(\xi, t) \sim \int_{-1}^1 dx \frac{GPD(x, \xi, t)}{x - \xi}. \quad (5)$$

Experimentally, the DVCS amplitude interferes with the Bethe-Heitler process (Fig.2) and

$$\sigma = |\mathcal{A}_{BH} + \mathcal{A}_{DVCS}|^2 = |\mathcal{A}_{BH}|^2 + |\mathcal{A}_{DVCS}|^2 + 2\Re \{ \mathcal{A}_{BH} \mathcal{A}_{DVCS}^* \} \quad (6)$$

while  $\Im \mathcal{A}_{DVCS}(\xi, t)$  can be separated using the beam spin asymmetry, and  $\Re \mathcal{A}_{DVCS}(\xi, t)$  from the angular dependence, the beam charge asymmetry ( $e^+$  v.  $e^-$  or  $\mu^+$  v.  $\mu^-$ ) can be used as an independent means to isolate the real part of the DVCS amplitude: the

interference terms between BH and DVCS depends on the sign of the lepton charge! For more details regarding the measurement of the DVCS amplitude, see Ref. [1] and references therein.

## PHYSICS OF GPDs

One of the reasons GPDs attracted interest is that the  $2^{nd}$  moment of  $H^q + E^q$  can be identified with the angular momentum (spin plus orbital) carried by quarks with flavor  $q$  in a nucleon with spin polarization  $\vec{S}$  [2]

$$\langle \vec{J}^q \rangle \equiv \langle q^\dagger \vec{\Sigma} q \rangle + \langle q^\dagger \vec{r} \times (\vec{p} - q\vec{A}) q \rangle = \vec{S} \int_{-1}^1 dx x [H^q(x, \xi, 0) + E^q(x, \xi, 0)]. \quad (7)$$

Note that while the  $\xi$ -dependence (term  $\propto \xi^2$ ) on the r.h.s. cancels between  $H^q(x, \xi, 0)$  and  $E^q(x, \xi, 0)$ , those integrals must be performed at the same fixed value of  $\xi$ .

Another important aspect of GPDs is their connection with *impact parameter dependent parton distributions*  $q(x, \mathbf{b}_\perp)$ . The latter are defined similar to the usual PDFs, but at distance  $\mathbf{b}_\perp$  relative to the *transverse center of longitudinal momentum*  $\mathbf{R}_\perp \equiv \sum_{i \in q, g} x_i \mathbf{b}_{\perp, i}$ . The Fourier transform of GPDs for  $\xi = 0$  yields these impact parameter dependent PDFs. For example, the two-dimensional Fourier transform of  $H^q(x, 0, -\Delta_\perp^2)$  yields the distribution of unpolarized quarks in an unpolarized or longitudinally polarized hadron [3]

$$q(x, \mathbf{b}_\perp) = \int \frac{d^2 \Delta_\perp}{(2\pi)^2} H^q(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp}. \quad (8)$$

The transverse gradients of the Fourier transform of  $E^q(x, 0, -\Delta_\perp^2)$  describes the transverse deformation of the distribution of unpolarized quarks in a transversely polarized nucleon [4]. For more details, see Ref. [1] and references therein.

Unfortunately, the DVCS amplitude is mostly sensitive to the regime  $x \approx \xi$ . In fact, the imaginary part is sensitive to  $x = \xi$  only (4), while the convolution integral for the real part (5) is dominated by the vicinity of  $x \approx \xi$ .

## INFORMATION CONTENT OF THE DVCS AMPLITUDE

In the following we will focus on charge even GPDs  $GPD^+(x, \xi, t) \equiv GPD(x, \xi, t) - GPD(-x, \xi, t)$ . Furthermore, the  $Q^2$  dependence will not be shown explicitly although all terms depend on  $Q^2$ .

Using dispersion relations one can show that [5]

$$\Re \mathcal{A}_{DVCS} \sim \int_{-1}^1 dx \frac{GPD^+(x, \xi, t)}{x - \xi} = \int_{-1}^1 dx \frac{GPD^+(x, x, t)}{x - \xi} + \Delta(t), \quad (9)$$

where  $\Delta(t)$  is the D-form factor [6]. This remarkable relation also follows from polynomiality [7] and implies that the information content of the DVCS amplitude (at fixed  $Q^2$ ) can be condensed to GPDs along the diagonal  $x = \xi$  plus the D-form factor. It

should be emphasized that  $\Re \mathcal{A}_{DVCS}(\xi, t)$  still adds more information to  $\Im \mathcal{A}_{DVCS}(\xi, t)$  than just  $\Delta(t)$ , since, for fixed  $t$  not the whole range  $0 < \xi < 1$  is accessible (at very low  $\xi$  the Bjorken limit may not yet have been reached and high  $\xi$  is inaccessible since  $t = -\frac{4\xi^2 M^2 + \Delta^2}{1 - \xi^2}$ ), while the above integrals extend from  $\xi = 0$  to  $\xi = 1$ . Nevertheless, (9) suggests to fit parameterizations for  $GPD(\xi, \xi, t)$  and  $D(t)$  to DVCS data rather than attempting to constrain parameters for  $GPD(x, \xi, t)$  over the whole  $x - \xi$  range.  $GPD(\xi, \xi, t)$  and  $\Delta(t)$  could then be used as an interface between experimental data and models. In fact, due to (9), any model/parameterization of GPDs satisfying polynomiality that fits both  $\Re \mathcal{A}_{DVCS}$  and  $\Im \mathcal{A}_{DVCS}$ , would also fit  $GPD(\xi, \xi, t)$  and  $\Delta(t)$  and vice versa. Moreover, fitting GPD-models to DVCS-data is not unique. For example, one can always fit DVCS data with the ansatz

$$H(x, \xi, t) = H_{DD}(x, \xi, t) + \Theta(|\xi| - |x|)D\left(\frac{x}{\xi}, t\right) \quad (10)$$

where for the ‘double distribution’ [9] one makes the specific ansatz  $H_{DD}(x, \xi, t) = H(x, x, t)$ , and  $D(z, t)$  being an arbitrary function that satisfies  $\int_{-1}^1 \frac{dz}{z-1} D(z, t) = \Delta(t)$ .

With the information from DVCS reduced to GPDs along the diagonal plus the  $D$ -form factor, the question arises what one can learn from this information. GPDs along the diagonal have been discussed e.g. in Ref. [8].

One very interesting observable that one can extract from this information is the  $\frac{1}{x}$ -moment of GPDs for  $\xi = 0$ . Provided  $t$  is large enough that the limit exists, one can take the  $\xi \rightarrow 0$  limit in the remarkable relation on the r.h.s. of (9), yielding

$$\int_{-1}^1 dx \frac{GPD^+(x, 0, t)}{x} = \int_{-1}^1 dx \frac{GPD^+(x, x, t)}{x} + \Delta(t). \quad (11)$$

From the GPD along the diagonal plus the  $D$ -form factor one can thus obtain the same  $\frac{1}{x}$ -moment of GPDs with  $\xi = 0$  that also enters the wide angle Compton scattering (WACS) [10] amplitudes. The main advantage of the approach outlined here compared to WACS is that using DVCS and (11), one can access  $\int_{-1}^1 dx \frac{GPD^+(x, 0, t)}{x}$  in a regime where  $M^2 < -t \ll Q^2$ , where the clear separation of scales facilitates the interpretation in terms of factorization.

Knowledge of  $\int_{-1}^1 dx \frac{GPD^+(x, 0, t)}{x}$  would be very valuable for understanding the physics of form factors at large  $-t$ : since antiquarks and sea quarks are not expected to play a significant role at large  $-t$ , a flavor separation of the  $u$  and  $d$  contributions should be possible using proton and neutron data only. Comparing  $\int_{-1}^1 dx \frac{GPD^+(x, 0, t)}{x}$  for each quark flavor with flavor separated form factor data should help understand what type of quarks dominate those form factors at large momentum transfer. For example, if the ratio between  $\int_{-1}^1 dx \frac{GPD^+(x, 0, t)}{x}$  and the corresponding form factor approaches 1 at large  $-t$  then that form factor would be dominated by quarks at  $x \rightarrow 1$  in that limit, while if that ratio turns out to be significantly greater than 1 for large  $-t$  then the corresponding form factor would be dominated by quarks carrying intermediate  $x$ .

The interpretation of the  $D$ -form factor remains obscure. While an interpretation of the  $z$ -moment of the  $D$ -term has been provided in terms of the stress-tensor [6], the  $D$ -form factor is the  $\frac{1}{z}$  moment of the  $D$ -term. This  $\frac{1}{z}$  moment also contributes a  $\delta(x)$ -type

contribution to  $\Re \mathcal{A}_{DVCS}$  in the  $\xi \rightarrow 0$  limit, since  $H(x, \xi, t) = H_{DD}(x, \xi, t) + \Theta(|\xi| - |x|)D\left(\frac{x}{\xi}\right)$  on the one hand, i.e. for  $x \neq 0$  one finds  $\lim_{\xi \rightarrow 0} H(x, \xi, t) = H_{DD}(x, \xi, t)$ , but on the other hand  $\lim_{\xi \rightarrow 0} \Re \mathcal{A}_{DVCS}(\xi, t) = \int_{-1}^1 \frac{dx}{x} H(x, 0, t) = \int_{-1}^1 \frac{dx}{x} H_{DD}(x, 0, t) = \Delta(t)$ , i.e.

$$\frac{H(x, 0, t) - H_{DD}(x, 0, t)}{x} = \delta(x)\Delta(t). \quad (12)$$

## **$Q^2$ -EVOLUTION**

One possibility to resolve the non-uniqueness of GPD extraction from DVCS data is the study of ‘double DVCS’ (DDVCS), where the photon on the ‘final state’ is also virtual, i.e. rather than producing a real photon, for example a lepton pair is produced. An alternative to this difficult process might be using the  $Q^2$  evolution of GPDs. To illustrate this point, imagine in the study of ordinary PDFs one were able to measure PDFs only at one value of  $x$ , but over a wide range of  $Q^2$ . Since the DGLAP evolution equations that govern the  $Q^2$  dependence of PDFs are known, one could thus still (partly) reconstruct the PDFs over a broad  $x$  range from this information.

In the context of GPDs, one also knows the evolution equations. Therefore, even if one can access GPDs only along the line  $x = \xi$ , since the  $x$ -distribution changes under  $Q^2$ , and since the Kernels that govern this evolution are also known, one can use the  $Q^2$  dependence to help disentangle the  $x$ -dependence.

Of course, for this procedure to work, even the lowest values of  $Q^2$  used in such a fit would have to be high enough to ensure that higher twist effects — which are usually not accounted for in evolution equations — are certain to be absent.

## **ACKNOWLEDGMENTS**

This work was supported by the DOE under grant numbers DE-FG03-95ER40965 and DE-AC05-06OR23177 (under which Jefferson Science Associates, LLC, operates Jefferson Lab).

## **REFERENCES**

1. M. Burkardt, A. Miller, and W.-D. Nowak, arXiv:08122208
2. Phys. Rev. Lett. **78**, 610 (1997).
3. M. Burkardt, Phys. Rev. D **62**, 071503 (2000), Erratum-ibid. D **66**, 119903 (2002).
4. M. Burkardt, Int. J. Mod. Phys. A **18**, 173 (2003).
5. M. Diehl and D.Y. Ivanov, Eur. Phys. J. C **52**, 919 (2007).
6. M.V. Polyakov and C. Weiss, Phys. Rev. D **60**, 114017 (1999).
7. I.V. Anikin and O.V. Teryaev, Phys. Rev. D **76**, 056007 (2007).
8. M. Burkardt, arXiv:0711.1881
9. A.V. Radyushkin, Phys. Rev. D **56**, 5524 (1997).
10. A.V. Radyushkin, Phys. Rev. D **58**, 114008 (1998).